arXiv:0809.2892v2 [hep-ph] 23 Oct 2008

Bose-Einstein condensation in dense quark matter

Jens O. Andersen

Department of Physics, Norwegian University of Science and Technology, N-7491 Trondheim, Norway

---

Abstract

We consider the problem of Bose condensation of charged pions in QCD at finite isospin chemical potential μ_I using the $O(4)$-symmetric linear sigma model as an effective field theory for two-flavor QCD. Using the 2PI $1/N$-expansion, we determine the quasiparticle masses as well as the pion and chiral condensates as a function of the temperature and isospin chemical potential in the chiral limit and at the physical point. The calculations show that there is a competition between the condensates. At $T = 0$, Bose condensation takes place for chemical potentials larger than m_π. In the chiral limit, the chiral condensate vanishes for any finite value of μ_I.

Key words: Bose-Einstein condensation, dense quark matter
PACS: 11.10 Wx, 11.15 Bt, 12.38.Mh, 21.65.Qr

---

1. Introduction

Condensation of bosons occurs in various contexts of high-density QCD. In the color-flavor-locked (CFL) phase of high-baryon density QCD, the $SU(3)_c \times SU(3)_L \times SU(3)_R \times U(1)_B$ is broken down to $SU(3)_{c+L+R}$ and the breaking of the global symmetries gives rise to (pesudo)-Goldstone bosons. The lightest of the excitations are expected to be the neutral and charged kaons. If the chemical potentials are large enough and the temperature low enough, the kaons will condense. Kaon condensation in the CFL phase is discussed e.g. in Ref. [1] and also in the talk given by Andreas Schmitt at this conference [2].

Another interesting system in this context is QCD at finite isospin chemical potential μ_I. If the isospin chemical potential is large enough and the temperature sufficiently low, the charged pions will condense. In contrast to QCD at finite baryon chemical potential, QCD at finite isospin chemical potential is accessible on the lattice since there is no fermion sign problem. Lattice simulations [3] indicate that there is a deconfinement transition of pions at high temperature and low density, and Bose-Einstein condensation of charged pions at high isospin density and low temperature. Various aspects of

Preprint submitted to Elsevier *23 October 2008*

pion condensation have been investigated using chiral perturbation theory [4], (Polyakov loop)NJL models [5,6,7], and linear sigma models [6,8,9].

In this talk, I would like to present some results for Bose condensation of charged pions using the linear sigma model as a low-energy effective for two-flavor QCD. The talk is based on Refs. [8,9] and part of the work is done in collaboration with Tomas Brauner.

2. Interacting Bose gas and 2PI $1/N$-expansion

The Euclidean Lagrangian for a Bose gas with N species of massive charged scalars is

$$\mathcal{L} = (\partial_\mu \Phi_i^\dagger)(\partial_\mu \Phi_i) - \frac{H}{\sqrt{2}}\left(\Phi_1 + \Phi_1^\dagger\right) + m^2 \Phi_i^\dagger \Phi_i + \frac{\lambda}{2N}\left(\Phi_i^\dagger \Phi_i\right)^2 \;, \tag{1}$$

where $i = 1, 2, ..., N$ and $\Phi_i = (\phi_{2i-1} + i\phi_{2i})/\sqrt{2}$ is a complex field. A nonzero value of H allows for explicit symmetry breaking and finite pion masses in the vacuum. The incorporation of a conserved charge Q_i is done by making the substitution $\partial_0 \Phi_i \to (\partial_0 - \mu_i)\,\Phi_i$ in the Lagrangian (1). We introduce nonzero expectation values ϕ_0 and ρ for ϕ_1 and ϕ_2, respectively, to allow for a chiral condensate and a charged pion condensate. We then write the complex field Φ_1 as $\Phi_1 = \frac{1}{\sqrt{2}}(\phi_0 + i\rho_0 + \phi_1 + i\phi_2)$, where ϕ_1 and ϕ_2 are quantum fields. The inverse tree-level propagator D_0^{-1} reads

$$D_0^{-1} = \begin{pmatrix} \omega_n^2 + p^2 + m_1^2 & \frac{\lambda}{N}\phi_0\rho_0 & 0 & 0 \\ \frac{\lambda}{N}\phi_0\rho_0 & \omega_n^2 + p^2 + m_2^2 & -2\mu_I\omega_n & 0 \\ 0 & 2\mu_I\omega_n & \omega_n^2 + p^2 + m_3^2 & 0 \\ 0 & 0 & 0 & \omega_n^2 + p^2 + m_4^2 \end{pmatrix} \;, \tag{2}$$

where $p = |\mathbf{p}|$, $\omega_n = 2n\pi T$ is the nth Matsubara frequency, and the treel-level masses are

$$m_1^2 = m^2 + \frac{3\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2 \;, \quad m_2^2 = -\mu_I^2 + m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{3\lambda}{2N}\rho_0^2 \;, \tag{3}$$

$$m_3^2 = -\mu_I^2 + m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2 \;, \quad m_4^2 = m^2 + \frac{\lambda}{2N}\phi_0^2 + \frac{\lambda}{2N}\rho_0^2 \;. \tag{4}$$

We have introduced a chemical potential μ_I for the third component of the isospin charge, which corresponds to the current density $j^\mu \sim \phi_2\partial^\mu\phi_3 - \phi_3\partial^\mu\phi_2$. For nonzero μ_I, the charged pions $\pi^\pm$ are linear combinations of ϕ_2 and ϕ_3.

After symmetry breaking, the 2PI effective potential can be written as

$$\begin{aligned}\Gamma[\phi_0, \rho_0, D] = {}& \frac{1}{2}m^2\left(\phi_0^2 + \rho_0^2\right) + \frac{\lambda}{8N}\left(\phi_0^2 + \rho_0^2\right)^2 - \frac{1}{2}\mu_I^2\rho_0^2 - H\phi_0 + \frac{1}{2}\mathrm{Tr}\ln D^{-1} \\ & + \frac{1}{2}\mathrm{Tr} D_0^{-1} D + \Phi[D] \;,\end{aligned} \tag{5}$$

where D is the exact propagator and $\Phi[D]$ is the sum of all two-particle irreducible vacuum diagrams. The contributions to $\Phi[D]$ can be classified according to which order in $1/N$ they contribute. The leading-order contribution is the double-bubble diagram where the $2N - 3$ particles with tree-level masses m_4 are propagating in the loops.

The self-energy in the large-N limit, $\Pi_{\rm LO}$, is given by the tadpole diagram and is independent of momentum. The inverse propagator can then be written as $D^{-1}=p^2+\omega_n^2+m_4^2+\Pi_{\rm LO}$. The term $m_4^2+\Pi_{\rm LO}$ is thus a local mass term, which we denote by M^2. The exact propagator D satisfies the equation $\delta\Gamma/\delta D=0$. In the present case, it reduces to a local gap equation for M^2:

$$M^2=m_4^2+\lambda T\sum_{\omega_n=2\pi nT}\int_p\frac{d^dq}{(2\pi)^d}\frac{1}{\omega_n^2+q^2+M^2}\;. \tag{6}$$

The fields ϕ_0 and ρ_0 satisfy the stationarity conditions $\delta\Gamma/\delta\phi_0=0$ and $\delta\Gamma/\delta\rho_0=0$, i.e.

$$0=m^2\phi_0-H+\frac{\lambda}{2N}\phi_0\left(\phi_0^2+\rho_0^2\right)+\lambda\phi_0T\sum_{\omega_n}\int_p\frac{d^dq}{(2\pi)^d}\frac{1}{\omega_n^2+q^2+M^2}\;, \tag{7}$$

$$0=\left(m^2-\mu_I^2\right)\rho_0+\frac{\lambda}{2N}\rho_0\left(\phi_0^2+\rho_0^2\right)+\lambda\rho_0T\sum_{\omega_n}\int_p\frac{d^dq}{(2\pi)^d}\frac{1}{\omega_n^2+q^2+M^2}\;. \tag{8}$$

The gap equations require renormalization and details can be found in e.g. [8]. The diagrammatic interpretation of the gap equations is that they sum up all daisy and superdaisy diagrams. The value of M is found by solving the renormalized versions of the equations (6), (7), and (8). The quasiparticle masses are then found by solving the equation $\det D(\omega,p=0)=0$, where the full propagator is obtained from the tree-level propagator Eq. (2) by the replacement $m_i^2\rightarrow M_i^2=m_i^2+\Pi_{\rm LO}$.

3. Results

In Fig. 1, we show the quasiparticle masses as a function of isospin chemical potential for $T=0$ and at the physical point. The three pions are degenerate in the vacuum ($\mu_I=0$) and two of them become degenerate in the limit $\mu_I\rightarrow\infty$. Also notice that π^+ becomes massless at the the critical value of $\mu_I=\mu_I^c=m_\pi=139$ MeV, which signals the onset of Bose condensation. This is a Goldstone bosons arising from the breaking of the $O(2)$-symmetry. In Fig. 1, we also show the pion condensate as a function of T and μ_I. Notice the onset of Bose condensation for $\mu_I=m_\pi$ at $T=0$. The phase transition is of second order in accordance with universality arguments.



Fig. 1. Left: quasiparticle masses at $T=0$ at the physical point as a function of μ_I. Right: pion condensate at the physical point as a function of T and μ_I.

4. Summary and outlook

In this talk I have presented some results for Bose-Einstein condenstion of charged pions in QCD at finite isospin chemical potential. Using the 2PI $1/N$-expansion, I have calculated the quasiparticle masses as well as the chiral condensate and the pion condensate as a function of T and μ_I. Our large-N approximation is of mean-field type and one would like to go beyond mean-field theory by including the next-to-leading order in the $1/N$-expansion. The appearance of massless excitations is in accordance with Goldstone's theorem, which is respected by the 2PI $1/N$-expansion.

If Bose-condensation takes place in a star, the system must be globally electric as well color neutral, otherwise one has to pay an enormous energy penalty. In the context of kaon condensation, one would include electric charge neutrality by solving the equation $\partial\Gamma/\partial\mu_Q = 0$, where μ_Q is the electric chemical potential. The problem of kaon condensation in the CFL phase was investigated in detail using an $O(2) \times O(2)$ effective scalar in the Hartree approximation in the paper by Alford, Braby, and Schmitt [1], but without adressing renormalization issues. Work in this direction is in progress [10]. See also Ref. [11] for renormalization of the 2PI Hartree approximation.

Acknowledgments: Part of the work presented in this talk was done in collaboration with Tomas Brauner. The author would like to thank the organizers of SEWM2008 for a stimulating meeting.

References

- [1] M. Buballa, Phys. Lett. **B609**, 57 (2005); M. G. Alford, M. Braby, and A. Schmitt, J. Phys. **G**: nucl. Part. Phys. **35**, 025002 (2008); M. G. Alford, A. Schmitt, K. Rajagopal, and T. Schafer, arXiv:0709.4635, to appear in Rev. Mod. Phys.
- [2] A. Schmitt, these Proceedings.
- [3] J. B. Kogut and D. K. Sinclair, Phys. Rev. D **64** 014508 (2002); ibid D **66** 34505 (2002); S. Gupta, hep-lat/0202005; P. de Forcrand, M. A. Stephanov, U. Wenger, PoS LAT2007, (237) 2007.
- [4] K. Splittorff, D. T. Son and M. Stephanov, Phys. Rev. D **64** 016003 (2001). J. B. Kogut and D. Toublan, Phys. Rev. D **64** 034007 (2001); M. Loewe and Villavicencio, Phys. Rev. D **67**, 074034 (2003); ibid D **70**, 074005 (2004); ibid D **71**, 094001 (2005).
- [5] A. Barducci, R. Casalbuoni, G. Pettini, and L. Ravagli, Phys. Rev. D **69**, 096004 (2004): D. Ebert and K. G. Klimenko, Eur. Phys. J. **C46**, 771 (2006); J. O. Andersen and L. Kyllingstad, hep-ph/0701033; D. Zablocki, D. Blaschke, and R. Anglani, arXiv:0805.2687. H. Abuki, R. Anglani, R. Gatto, M. Pellicoro, and M. Ruggieri, arXiv:0809.2658.
- [6] L. He, M. Jin, and P. Zhuang, Phys.Rev. D **71**, 116001 (2005).
- [7] S. Mukherjee, M. G. Mustafa, and R. Ray, Phys. Rev. D **75**, 094015 (2007); C. H. Abuki, M. Ciminale, R. Gatto, G. Nardulli, and M. Ruggieri, Phys. Rev. D **77**, 014040 (2008).
- [8] J. O. Andersen, Phys. Rev. D, **75**, 065011 (2007).
- [9] J. O. Andersen and T. Brauner, Phys. Rev. D, **78**, 014030 (2008).
- [10] J. O. Andersen and L. Leganger, in preparation.
- [11] J. Berges, Sz. Borsanyi, Urko Reinosa, and J. Serreau, Annals Phys. **320**, 344 (2005); G. Fejos, A. Patkos, and Zs. Szep, Nucl. Phys. **A803**, 115 (2008); A. Patkos and Zs. Szep, arXiv:0809.3551 (these Proceedings); A. Jakovac, arXiv:0808.1800 and these Proceedings.